

Hb S/Hb Lepore With Mild Sickling Symptoms: A Hemoglobin Variant With Mostly δ -Chain Sequences Ameliorates Sickle-Cell Disease

V.F. Fairbanks,^{1*} D.J. McCormick,² K.S. Kubik,¹ W.N. Rezuze,³ D. Black,⁴ M.S. Ochaney,⁵ and D. Schwartz⁵

¹Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, Minnesota

²Department of Biochemistry and Molecular Biology, Mayo Clinic, Rochester, Minnesota

³Department of Pathology and Laboratory Medicine, Hartford Hospital, Hartford, Connecticut

⁴Department of Pediatrics, Burdorf Health Center, University of Connecticut-Mount Sinai Pediatric Module, Hartford, Connecticut

⁵Department of Internal Medicine, North Arundel Hospital, Glen Burnie, Maryland

Three cases are reported of Hb S/Hb Lepore combination with very mild sickling manifestations. The presence of a non- α -chain variant with a high proportion of δ chain sequences, including 22 ala, appears to ameliorate sickle-cell disease. Efforts to increase the proportion of Hb A₂ may be beneficial in sickle-cell disease. *Am. J. Hematol.* 54:164–165

© 1997 Wiley-Liss, Inc.

Key words: sickle-cell disease; Hb Lepore; Hb S; delta chain; Hb A₂

INTRODUCTION

Attempts to treat sickle-cell disease by increasing the proportion of Hb A₂ may have deleterious effects, including exacerbation of anemia [1]. However, constructs with the β -globin substitution 22-glu→ala inhibit Hb S polymerization [1]; Hb A₂ has 22 ala in the δ chain. Hb G-Coushatta is β 22 glu→ala. Hb Lepore-Boston has mostly δ chain sequences, including 22 ala. It would be informative to examine sickle-cell disease severity in patients with the combination Hb S/G-Coushatta or Hb S/Lepore-Boston, since such experiments of nature might answer questions posed by recent reports [1,2].

We have not encountered a case of Hb S/Hb G-Coushatta, nor are we aware of any reported cases. In such a case, the proportion of Hb S should approximate that in Hb S trait, and not cause serious sickling. However, the Hb S/Hb Lepore combination may have enough Hb S to cause serious sickling, particularly if an excess of δ chains were to have harmful consequences, as has been asserted. Few such cases have been previously reported [3–5]. In 4 of 7 previously reported cases of HbS/Hb Lepore, sickling manifestations were mild. In 3 other cases, manifestations were severe, although Hb concentration in these cases ranged from 8–9 g/dl. Three of 4 cases with mild sickling features were age 57, 60, and 76 years.

We describe 3 cases of Hb S/Hb Lepore, with relatively mild manifestations.

Case 1 is a 23-year-old African-American male who has been repeatedly admitted to hospital for treatment of pain in his back or legs, but with no other manifestations of sickle-cell disease. Physical examinations have been unremarkable.

Case 2, the 26-year-old sister of case 1, has recurrent arthralgias and acute decline in Hb concentration associated with acute infections, but has rarely been transfused.

Case 3 is a 16-year-old Jamaican black male who has experienced a few episodes of knee or elbow pain. He has no other manifestations of sickle-cell disease. His physical examination was unremarkable.

In the specimens from cases 1 and 2, identification of Hb Lepore was made on the basis of its characteristic electrophoretic mobility and by isoelectric focusing and HPLC with a weak cationic exchange resin column. In the specimen from case 3, Hb Lepore was not suspected initially because of its high proportion. An abnormal glo-

*Correspondence to: V.F. Fairbanks, Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN 55905.

Received for publication 23 May 1996; Accepted 18 June 1996.

TABLE I. Laboratory Data

Case no.	Hb conc. (g/dl)	RBC $\times 10^{12}/l$	MCV (fl)	Hemoglobin components (%) ^a				
				A	F	A ₂	S	Lepore
1	9.9	3.93	77.8	0	8	2	77	12
2	10.4	4.13	74.6	0	7	4	78	11
3	11.9	5.61	66.5	0	5	3	72	20

^aProportions of Hb components have been rounded to nearest whole integer.

bin chain, not β^S , was separated by HPLC, and its tryptic peptides were examined. Peptides 10 and 11 had δ sequences; peptide 12 had a β sequence; these results were consistent with Hb Lepore-Boston.

The microcytosis in these cases was due to Hb Lepore trait. The mildness of the symptoms suggests that the presence of a Hb variant with δ -chain sequences has no adverse effect when present together with high proportions of Hb S, but rather ameliorates sickle-cell disease. The high proportion of Hb Lepore in case 2 supports this conclusion.

Nagel et al. [1] indicate that human δ -globin chains bind avidly to mouse red-cell membranes, causing severe erythrocyte deformity and hemolysis. Their observations, in conflict with the mildness of the HbS/Hb Lepore sickling disorder, may be explained thus: either 1) human δ -globin chains may not have the same deleterious effect on human erythrocyte membranes, or else 2) the short segment of the human δ -globin chain, beyond the point

at which crossing-over occurs in Hb Lepore-Boston, may confer the deleterious effects observed by Nagel et al. [1], but the long segment containing δ -chain sequences, from amino-acid residue 1–87, does not.

REFERENCES

1. Nagel RL, Sharma A, Kumar R, Fabry ME: Severe red cell abnormalities in transgenic mice expressing high levels of normal human delta chains. *Blood [Suppl]* 86:251, 1995.
2. Reilly MP, McCune S, Townes TM, Asakura T: Characterization of an anti-sickling hemoglobin for gene therapy of sickle cell disease. *Blood [Suppl]* 86:417, 1995.
3. Ahern VN, Aarons GH, Jones RT, Brimhall B: Hemoglobin Lepore Washington in two Jamaican families: Interaction with beta chain variants. *Blood* 40:246–256, 1972.
4. Stamatoyannopoulos G, Fessas P: Observations on hemoglobin “Pylos”: The hemoglobin Pylos-hemoglobin S combination. *J Lab Clin Med* 62:193–200, 1963.
5. Silvestroni E, Bianco I, Baglioni C: Interaction of hemoglobin Lepore with sickle cell trait and microcythemia (thalassemia) in a Southern Italian family. *Blood* 25:457–469, 1965.